

Antisymmetry in Strangeness -1 and -2 Three-Baryon Systems

W. Glöckle

Institut für theoretische Physik II, Ruhr-Universität Bochum, D-4478 Bochum, Germany

K. Miyagawa

Applied Physics, Okayama University of Science, 1-1 Ridai-cho, Okayama 700, Japan

(February 8, 2008)

Abstract

Using the generalized Pauli principle by adding particle labels to the usual space and spin labels a symmetric Hamiltonian and a corresponding anti-symmetric wavefunction is constructed for systems of three baryons in the strangeness sectors $S = -1$ and -2 . Applications are the $\Xi NN - \Lambda\Lambda N$ and $NN\Lambda - NN\Sigma$ systems. Minimal sets of generalized coupled Faddeev equations for breakup and rearrangement operators as well as (possible) bound states are derived which have the ordinary Pauli principle among identical particles built in. The equations found confirm our previous sets of coupled Faddeev equations for those systems whose derivation was carried through for distinguishable particles and not using the generalized Pauli principle.

Typeset using REVTeX

I. INTRODUCTION

In the three-baryon system ΞNN a strong transition occurs into YYN where Y is either a Λ or a Σ . The nucleon present in YYN can be either one contained in ΞNN . Clearly the ΞNN part of the total wavefunction has to be antisymmetrical under exchange of the two nucleons. So the question arises how is that reflected in the second part of the wavefunction belonging to YYN . Similarly the second part has to be antisymmetrical under exchange of two identical hyperons and one has to ask how is that related to the first part of the wavefunction. There is also the possibility that YYN appears as $\Lambda\Sigma N$. Those questions (with the neglect of the Σ -admixture) have been answered in [1] starting from a second quantisation picture of the system. The final aim in [1] was to write down the set of coupled Faddeev equations with the smallest number of independent Faddeev amplitudes. The argumentation turned out to be rather involved and we would like to present a conceptually and algebraically simpler derivation of the same set of coupled equations. Since it is simpler we will now also include the possibility of a Σ -admixture.

The Ξd system has recently received attention [2] in that it might be a source of information on the $\Lambda\Lambda$ scattering length after it has switched to the unbound $\Lambda\Lambda N$ system. In that evaluation of those two coupled three-body systems the AGS equations [3] for distinguishable particles have been used and the free initial and final states have been antisymmetrised separately. Since "clean" experimental information on hyperon-hyperon forces are still very rare due to the lack of hyperon beams and targets the proposals to extract information on YN and YY forces from final state interactions have great importance and therefore a clearly justified formalism is in order.

Also under the present circumstances of having no direct access to YN scattering light hypernuclei play an equally important role, since they can be solved rigorously for baryon-baryon forces in all their complexities and thus provide an important test for those forces. First steps have been undertaken in [4] for the hypertriton and in [5] for ${}^4_{\Lambda}\text{He}$ and ${}^4_{\Lambda}\text{H}$.

With respect to investigations of final state interactions among hyperons and nucleons

the photon induced K^+Y production on the deuteron has been evaluated recently [6] again using modern YN forces which correctly bind the hypertriton. The corresponding electron induced process on the deuteron has been measured recently [7] and a first calculation appeared [8]. Certainly more work, both experimentally and theoretically, will come up in the near future.

In section II we describe the formalism of antisymmetrisation for particles of different types and apply it to the relatively simple two-baryon system. The more complex situation for three baryons with strangeness -1 and -2 and the derivation of coupled Faddeev equations is worked out in section III. An application to various three-body breakup processes initiated by a Ξ particle is described in section IV. Generalised AGS equations for elastic and rearrangement scattering are described in section V. We summarize in section VI.

II. FORMALISM AND APPLICATION TO THE TWO-BARYON SYSTEM

The ΞN system can strongly convert into two identical baryons, $\Lambda\Lambda$ or $\Sigma\Sigma$. Clearly the parts of the wavefunction describing $\Lambda\Lambda$ or $\Sigma\Sigma$ are antisymmetrical under exchange of the two particles, whereas for the part of the wavefunction describing ΞN there is no symmetry requirement. Another possible appearance of that strangeness -2 system is $\Lambda\Sigma$ with distinguishable particles. In [1] we used the formalism of second quantisation to derive three coupled equations for the coupled ΞN and $\Lambda\Lambda$ systems, neglecting a possible Σ -admixture. One of the equations can be dropped at the expense of working with a non-symmetrical 2×2 potential matrix.

In the np system it has been known since the early days of nuclear physics that adding an isospin label to the particle or in other words extending the space of states by introducing isospin one can require that the wavefunction for the distinguishable particles n and p has to be antisymmetrical. The simple mechanism is that in isospin space there is a symmetrical and antisymmetrical basis which can be combined with the corresponding antisymmetrical and symmetrical wavefunction in ordinary and spin space. Therefore no restriction has been

introduced by requiring antisymmetry in the enlarged space. Connected to that antisymmetric wavefunction a symmetric Hamiltonian can be written down. This idea has been generalized also long time ago to SU(3) including strangeness on top of isospin. If the Hamiltonian severely breaks SU(3) symmetry due to significantly different masses and symmetry breaking force parameters the use of irreducible representations of SU(3) appears not to be of much help and a formally simpler approach can be used. Again we think this is known and we display it again for two particles to lay the ground for the more complex three-baryon system, where we want to find the way to the smallest set of coupled Faddeev equations.

Let us introduce states $|\alpha\beta\rangle$ with particle labels α or β denoting nucleons or hyperons. Thus $\alpha = \{N, \Lambda, \Sigma, \Xi\}$. Note $|\alpha\beta\rangle$ stands for $|\alpha\rangle_1 |\beta\rangle_2$ which means that particle 1 is of the type α and 2 is of type β . Thus the transposition P_{12} yields $P_{12} |\alpha\beta\rangle = |\beta\alpha\rangle$. The completeness relation in that particle space is obviously

$$\sum_{\alpha\beta} |\alpha\beta\rangle\langle\alpha\beta| = 1 \quad (2.1)$$

Since strangeness is a good quantum number the sum is restricted to run only over physically allowed states. For $S = -1$ these are $|N\Lambda\rangle$, $|\Lambda N\rangle$, $|N\Sigma\rangle$, and $|\Sigma N\rangle$. For $S = -2$ one has $|\Lambda\Lambda\rangle$, $|\Sigma\Sigma\rangle$, $|\Xi N\rangle$, $|N\Xi\rangle$, $|\Lambda\Sigma\rangle$ and $|\Sigma\Lambda\rangle$. Clearly the sum in Eq. (2.1) is symmetrical under exchange of the two particles. Applying that sum onto the corresponding two-body Hamiltonian from both sides one creates a fully symmetrical Hamiltonian

$$\begin{aligned} H &\equiv \sum_{\alpha\beta} \sum_{\alpha'\beta'} |\alpha\beta\rangle\langle\alpha\beta| H |\alpha'\beta'\rangle\langle\alpha'\beta'| \\ &= \sum_{\alpha\beta} |\alpha\beta\rangle \left(\frac{p_1^2}{2m_\alpha} + \frac{p_2^2}{2m_\beta} + m_\alpha + m_\beta \right) \langle\alpha\beta| \\ &\quad + \sum_{\alpha\beta} \sum_{\alpha'\beta'} |\alpha\beta\rangle V_{\alpha\beta,\alpha'\beta'} \langle\alpha'\beta'| \end{aligned} \quad (2.2)$$

Note that we allow for different masses m_α without destroying the symmetry of the kinetic energy. In the configuration or momentum space representation including (possible)

spin dependencies the potentials are denoted by $V_{\alpha\beta,\alpha'\beta'}(12)$. In general these potential functions are not symmetrical. But they have the property

$$V_{\beta\alpha,\beta'\alpha'}(12) = V_{\alpha\beta,\alpha'\beta'}(21) \quad (2.3)$$

Transpositions applied to potential functions act on the arguments (momenta and spins) and thus one has

$$P_{12}V_{\alpha\beta,\alpha'\beta'}(12) = V_{\alpha\beta,\alpha'\beta'}(21)P_{12} \quad (2.4)$$

They do not change the labels for the potential functions, which are just parameters of those functions.

To that symmetrical Hamiltonian belongs obviously a fully antisymmetrical two-baryon wavefunction:

$$|\Psi\rangle = \sum_{\alpha\beta} |\alpha\beta\rangle \langle\alpha\beta|\Psi\rangle \equiv \sum_{\alpha\beta} |\alpha\beta\rangle \psi_{\alpha\beta} \quad (2.5)$$

Like the potential functions the amplitudes $\psi_{\alpha\beta}$ live in spin and configuration or momentum space and corresponding representations will be denoted by $\psi_{\alpha\beta}(12)$. For $\alpha \neq \beta$ one introduces two-particle states

$$|\chi_{\alpha\beta}^a\rangle \equiv \frac{1}{\sqrt{2}}(|\alpha\beta\rangle - |\beta\alpha\rangle) \quad (2.6)$$

$$|\chi_{\alpha\beta}^s\rangle \equiv \frac{1}{\sqrt{2}}(|\alpha\beta\rangle + |\beta\alpha\rangle) \quad (2.7)$$

with obvious symmetry properties. In addition there is the symmetrical state

$$\chi_{\alpha\alpha}^s \equiv |\alpha\alpha\rangle \quad (2.8)$$

Then the total state Ψ can be written as

$$|\Psi\rangle = \sum_{\alpha} |\alpha\alpha\rangle \psi_{\alpha\alpha} + \sum_{\alpha \neq \beta} \frac{1}{\sqrt{2}} (|\chi_{\alpha\beta}^a\rangle + |\chi_{\alpha\beta}^s\rangle) \psi_{\alpha\beta} \quad (2.9)$$

Now we impose antisymmetry

$$P_{12}|\Psi\rangle = -|\Psi\rangle \quad (2.10)$$

which leads after some inspection to

$$P_{12}\psi_{\alpha\beta} = -\psi_{\beta\alpha} \quad (2.11)$$

Note P_{12} in Eq. (2.10) acts on the particle labels of all states, while P_{12} in Eq. (2.11) acts only on those for momentum(configuration) and spin states. In other words Eq. (2.11) leads to

$$\psi_{\alpha\beta}(21) = -\psi_{\beta\alpha}(12) \quad (2.12)$$

The antisymmetry of (2.9) is explicit if we keep (2.11) in mind and rewrite (2.9) as

$$\begin{aligned} |\Psi\rangle &= \sum_{\alpha} |\alpha\alpha\rangle \psi_{\alpha\alpha} + \sum_{\alpha < \beta} |\chi_{\alpha\beta}^a\rangle \frac{1}{\sqrt{2}} (\psi_{\alpha\beta} - \psi_{\beta\alpha}) \\ &\quad + \sum_{\alpha < \beta} |\chi_{\alpha\beta}^s\rangle \frac{1}{\sqrt{2}} (\psi_{\alpha\beta} + \psi_{\beta\alpha}) \\ &= \sum_{\alpha} |\alpha\alpha\rangle \psi_{\alpha\alpha} + \sum_{\alpha < \beta} |\chi_{\alpha\beta}^a\rangle \frac{1}{\sqrt{2}} (1 + P_{12}) \psi_{\alpha\beta} \\ &\quad + \sum_{\alpha < \beta} |\chi_{\alpha\beta}^s\rangle \frac{1}{\sqrt{2}} (1 - P_{12}) \psi_{\alpha\beta} \end{aligned} \quad (2.13)$$

Here we introduced a definite but arbitrary ordering of the particle labels.

Now projecting $H\Psi = E\Psi$ onto $\langle 12 | \langle \alpha \beta |$ where $\langle 12 |$ stands for momenta and spin states, one obtains

$$\begin{aligned} & \left(\frac{p_1^2}{2m_\alpha} + \frac{p_2^2}{2m_\beta} + m_\alpha + m_\beta \right) \psi_{\alpha\beta}(12) + \sum_{\alpha' \neq \beta'} V_{\alpha\beta, \alpha'\beta'}(12) \psi_{\alpha'\beta'}(12) \\ & + \sum_{\alpha'} V_{\alpha\beta, \alpha'\alpha'}(12) \psi_{\alpha'\alpha'}(12) = E \psi_{\alpha\beta}(12) \end{aligned} \quad (2.14)$$

Again introducing the arbitrary but fixed ordering into the different particle labels leads to

$$\begin{aligned} & \left(\frac{p_1^2}{2m_\alpha} + \frac{p_2^2}{2m_\beta} + m_\alpha + m_\beta \right) \psi_{\alpha\beta}(12) + \sum_{\alpha' < \beta'} V_{\alpha\beta, \alpha'\beta'}^{tot}(12) \psi_{\alpha'\beta'}(12) \\ & + \sum_{\alpha'} V_{\alpha\beta, \alpha'\alpha'}(12) \psi_{\alpha'\alpha'}(12) = E \psi_{\alpha\beta}(12) \end{aligned} \quad (2.15)$$

Using Eq. (2.11) we introduced for $\alpha' \neq \beta$ the "total"-potential leading from $\alpha\beta$ to $\alpha'\beta'$ which is the sum of a "direct" potential and an "exchange" potential

$$V_{\alpha\beta, \alpha'\beta'}^{tot}(12) \equiv V_{\alpha\beta, \alpha'\beta'}(12) - V_{\alpha\beta, \beta'\alpha'}(12) P_{12} \quad (2.16)$$

The application of Eq. (2.15) to $|\alpha\beta\rangle = \{ |N\Lambda\rangle, |N\Sigma\rangle \}$ is obvious and will not be written down. For $|\alpha\beta\rangle \equiv \{ |\Xi N\rangle, |\Lambda\Lambda\rangle, |\Sigma\Sigma\rangle, |\Lambda\Sigma\rangle \}$ we obtain the set of four coupled equations

$$\begin{aligned} & \left(\frac{p_1^2}{2m_\Xi} + \frac{p_2^2}{2m_N} + m_\Xi + m_N - E \right) \psi_{\Xi N}(12) + V_{\Xi N, \Xi N}(12) \psi_{\Xi N}(12) \\ & + V_{\Xi N, \Lambda\Sigma}^{tot}(12) \psi_{\Lambda\Sigma}(12) + V_{\Xi N, \Lambda\Lambda}(12) \psi_{\Lambda\Lambda}(12) + V_{\Xi N, \Sigma\Sigma}(12) \psi_{\Sigma\Sigma}(12) = 0 \end{aligned} \quad (2.17)$$

$$\left(\frac{p_1^2}{2m_\Lambda} + \frac{p_2^2}{2m_\Sigma} + m_\Lambda + m_\Sigma - E \right) \psi_{\Lambda\Sigma}(12) + V_{\Lambda\Sigma, \Lambda\Sigma}^{tot}(12) \psi_{\Lambda\Sigma}(12)$$

$$+V_{\Lambda\Sigma,\Xi N}^{tot}(12)\psi_{\Xi N}(12) + V_{\Lambda\Sigma,\Sigma\Sigma}(12)\psi_{\Sigma\Sigma}(12) = 0 \quad (2.18)$$

$$\begin{aligned} &(\frac{p_1^2}{2m_\Lambda} + \frac{p_2^2}{2m_\Lambda} + 2m_\Lambda - E)\psi_{\Lambda\Lambda}(12) + V_{\Lambda\Lambda,\Xi N}^{tot}(12)\psi_{\Xi N}(12) \\ &+ V_{\Lambda\Lambda,\Sigma\Sigma}(12)\psi_{\Sigma\Sigma}(12) + V_{\Lambda\Lambda,\Lambda\Lambda}(12)\psi_{\Lambda\Lambda}(12) = 0 \end{aligned} \quad (2.19)$$

$$\begin{aligned} &(\frac{p_1^2}{2m_\Sigma} + \frac{p_2^2}{2m_\Sigma} + 2m_\Sigma - E)\psi_{\Sigma\Sigma}(12) + V_{\Sigma\Sigma,\Xi N}^{tot}(12)\psi_{\Xi N}(12) + V_{\Sigma\Sigma,\Lambda\Sigma}^{tot}(12)\psi_{\Lambda\Sigma}(12) \\ &+ V_{\Sigma\Sigma,\Lambda\Lambda}(12)\psi_{\Lambda\Lambda}(12) + V_{\Sigma\Sigma,\Sigma\Sigma}(12)\psi_{\Sigma\Sigma}(12) = 0 \end{aligned} \quad (2.20)$$

This set has to be solved under the condition that $\psi_{\Lambda\Lambda}(12)$ and $\psi_{\Sigma\Sigma}(12)$ are antisymmetric, which imposes conditions on the wavefunction components with two different particle species. In [1] we found three coupled equations regarding however only the coupled $\Xi N - \Lambda\Lambda$ system. Dropping in Eqs. (2.17-20) the Σ state we have now two equations. The second of the three equations (2.15) from [1] has there been introduced in order to have a symmetrical potential matrix. But it results simply from the first equation in that set (2.15) by permuting the two particles and using the property (2.13) in [1] which is also given here as Eq. (2.12). The remaining two equations in [1] are identical to Eq. (2.17) and Eq. (2.19). The interaction $V_{\Xi N}$ in [1] includes the exchange term according to its construction and is the same as $V_{\Xi N,\Xi N}^{tot}$. As is seen from Eq. (2.13) the total Ψ contains both, symmetrical and antisymmetrical pieces, in the parts of the wavefunction referring to different particles.

We would like to add that in [1] unfortunately some errors sneaked in. After Eq. (2.3) it is said that the variables x or y include isospin quantum numbers, for instance. This is incorrect, x or y stand only for ordinary space and spin variables. Also the description of the examples following Eq. (2.22) is incorrectly given.

III. THREE-BARYON SYSTEMS

We start off by introducing three-particle states for total strangeness $S = -1$ and $S = -2$ systems. For instance in case of $S = -1$ they are $|\Lambda NN\rangle$, $|N\Lambda N\rangle$, $|N N\Lambda\rangle$, $|\Sigma NN\rangle$,

$|N\Sigma N\rangle$ and $|NN\Sigma\rangle$, whereas for $S = -2$ one has 15 states composed of N , Λ , Σ and Ξ species. A general state is denoted by $|\alpha\beta\gamma\rangle$ where α, β, γ run over the different species. Of course the sets of states contain only physically allowed combinations which conserve charge, strangeness and isospin. The completeness relation in particle space reads

$$\sum_{\alpha\beta\gamma} |\alpha\beta\gamma\rangle\langle\alpha\beta\gamma| = 1 \quad (3.1)$$

and the fully symmetrised Hamiltonian is

$$H = \sum_{\alpha\beta\gamma} \sum_{\alpha'\beta'\gamma'} |\alpha\beta\gamma\rangle\langle\alpha\beta\gamma| H |\alpha'\beta'\gamma'\rangle\langle\alpha'\beta'\gamma'| \quad (3.2)$$

The eigenstates Ψ of H are then totally antisymmetrical. In what follows we derive coupled Faddeev equations. The total state Ψ will first be decomposed into three Faddeev amplitudes, which due to the antisymmetry of Ψ can be written as [9]

$$\Psi = (1 + P)\psi \quad (3.3)$$

Here ψ is one of the three Faddeev amplitudes and $P = P_{12}P_{23} + P_{13}P_{23}$ is the sum of a cyclical and an anticyclical permutation of three objects. If ψ is antisymmetrical under exchange of two particles then Ψ given in Eq. (3.3) is fully antisymmetrical. That one Faddeev amplitude is based on one pair interaction contained in H , say between particles 2 and 3, and is defined as

$$\psi = G_0 V(23) \Psi \quad (3.4)$$

where G_0 is the free three-baryon propagator. Consequently it obeys the equation

$$\psi = G_0 V(23) \psi + G_0 V(23) P \psi \quad (3.5)$$

Since the symmetry condition for ψ is a simple one, namely to be antisymmetrical under exchange of particles 2 and 3, we can simply introduce a complete set of particle states which are antisymmetrical and symmetrical under exchange of particles 2 and 3:

$$|\chi_{\alpha\beta\gamma}^a\rangle \equiv \frac{1}{\sqrt{2}}(|\alpha\beta\gamma\rangle - |\alpha\gamma\beta\rangle) \quad \beta \neq \gamma \quad (3.6)$$

$$|\chi_{\alpha\beta\gamma}^s\rangle \equiv \frac{1}{\sqrt{2}}(|\alpha\beta\gamma\rangle + |\alpha\gamma\beta\rangle) \quad \beta \neq \gamma \quad (3.7)$$

$$|\chi_{\alpha\beta\beta}^s\rangle \equiv |\alpha\beta\beta\rangle \quad (3.8)$$

The general form

$$|\psi\rangle = \sum_{\alpha\beta\gamma} |\alpha\beta\gamma\rangle \langle \alpha\beta\gamma | \psi \rangle \equiv \sum_{\alpha\beta,\gamma} |\alpha\beta\gamma\rangle \psi_{\alpha\beta\gamma} \quad (3.9)$$

can then be rewritten as

$$|\psi\rangle = \sum_{\alpha\beta} |\chi_{\alpha\beta\beta}^s\rangle \psi_{\alpha\beta\beta} + \sum_{\alpha} \sum_{\beta \neq \gamma} \frac{1}{\sqrt{2}} (|\chi_{\alpha\beta\gamma}^a\rangle + |\chi_{\alpha\beta\gamma}^s\rangle) \psi_{\alpha\beta\gamma} \quad (3.10)$$

Imposing antisymmetry under exchange of particles 2 and 3 leads to

$$P_{23}\psi_{\alpha\beta\gamma} = -\psi_{\alpha\gamma\beta} \quad (3.11)$$

and after some inspection to

$$\begin{aligned} \psi &= \sum_{\alpha} \sum_{\beta < \gamma} |\chi_{\alpha\beta\gamma}^a\rangle \frac{1}{\sqrt{2}} (1 + P_{23}) \psi_{\alpha\beta\gamma} + \sum_{\alpha} \sum_{\beta < \gamma} |\chi_{\alpha\beta\gamma}^s\rangle \frac{1}{\sqrt{2}} (1 - P_{23}) \psi_{\alpha\beta\gamma} \\ &+ \sum_{\alpha\beta} |\chi_{\alpha\beta\beta}^s\rangle \psi_{\alpha\beta\beta} \end{aligned} \quad (3.12)$$

The set of coupled equations for $\psi_{\alpha\beta\gamma}$ is based on Eq. (3.5), which when projected reads

$$\psi_{\alpha\beta\gamma} = G_0^{\alpha\beta\gamma} (< \alpha\beta\gamma|V(23)|\psi > + < \alpha\beta\gamma|V(23)P|\psi >) \quad (3.13)$$

Clearly the free propagators $G_0^{\alpha\beta\gamma}$ are diagonal and depend on the particle species entering through their masses. The order $\alpha\beta\gamma$ corresponds of course to particles numbered 1,2 and 3. The first matrix element on the right hand side of Eq. (3.13) is easily evaluated as

$$\begin{aligned} & < \alpha\beta\gamma|V|\psi > \\ &= \sum_{\beta'\gamma'} \bar{\delta}_{\beta'\gamma'} V_{\beta\gamma,\beta'\gamma'}(23) < \alpha\beta'\gamma'|\psi > + \sum_{\beta'} V_{\beta\gamma,\beta'\beta'}(23) < \alpha\beta'\beta'|\psi > \\ &= \sum_{\beta' < \gamma'} (V_{\beta\gamma,\beta'\gamma'}(23) - V_{\beta\gamma,\gamma'\beta'}(23)P_{23})\psi_{\alpha\beta'\gamma'} + \sum_{\beta'} V_{\beta\gamma,\beta'\beta'}(23)\psi_{\alpha\beta'\beta'} \\ &\equiv \sum_{\beta' < \gamma'} V_{\beta\gamma,\beta'\gamma'}^{tot}(23)\psi_{\alpha\beta'\gamma'} + \sum_{\beta'} V_{\beta\gamma,\beta'\beta'}(23)\psi_{\alpha\beta'\beta'} \end{aligned} \quad (3.14)$$

Note we used the property (3.11) and introduced the convenient notation $\bar{\delta}_{\beta\gamma} \equiv 1 - \delta_{\beta\gamma}$.

In order to evaluate the second matrix element on the right hand side of Eq. (3.13) we need $P\psi$. Using (3.9) one finds after a simple algebra

$$< \alpha\beta\gamma|P|\psi > = P_{12}P_{23}\psi_{\beta\gamma\alpha} + P_{13}P_{23}\psi_{\gamma\alpha\beta} \quad (3.15)$$

This then leads easily to

$$\begin{aligned} < \alpha\beta\gamma|VP|\psi > = \sum_{\beta' < \gamma'} V_{\beta\gamma,\beta'\gamma'}^{tot}(23) \{P_{12}P_{23}\psi_{\beta'\gamma'\alpha} + P_{13}P_{23}\psi_{\gamma'\alpha\beta'}\} \\ &+ \sum_{\beta'} V_{\beta\gamma,\beta'\beta'}(23)(1 - P_{23})P_{13}P_{23}\psi_{\beta'\alpha\beta'} \end{aligned} \quad (3.16)$$

Here we used the property

$$P_{23}P\psi = -P\psi \quad (3.17)$$

which is evident from Eq. (3.12). Put together we end up with the following set of coupled equations for the amplitudes $\psi_{\alpha\beta\gamma}$

$$\begin{aligned}\psi_{\alpha\beta\gamma} = & G_0^{\alpha\beta\gamma} \left(\sum_{\beta' < \gamma'} V_{\beta\gamma, \beta'\gamma'}^{tot}(23) \psi_{\alpha\beta'\gamma'} + \sum_{\beta'} V_{\beta\gamma, \beta'\beta'}(23) \psi_{\alpha\beta'\beta'} \right) \\ & + G_0^{\alpha\beta\gamma} \left\{ \sum_{\beta' < \gamma'} V_{\beta\gamma, \beta'\gamma'}^{tot}(23) (P_{12}P_{23} \psi_{\beta'\gamma'\alpha} + P_{13}P_{23} \psi_{\gamma'\alpha\beta'}) \right. \\ & \left. + \sum_{\beta'} V_{\beta\gamma, \beta'\beta'}(23) (1 - P_{23}) P_{13}P_{23} \psi_{\beta'\alpha\beta'} \right\}\end{aligned}\quad (3.18)$$

We cast it into the following matrix form

$$\tilde{\psi} = \tilde{G}_0 \tilde{V} \tilde{\psi} + \tilde{G}_0 \tilde{V} \tilde{R} \quad (3.19)$$

where $\tilde{G}_0$ is diagonal and contains the various free propagators $G_0^{\alpha\beta\gamma}$. The matrix $\tilde{V}$ is composed of all non vanishing diagonal and transition potentials and $\tilde{R}$ of the $\psi_{\alpha\beta\gamma}$ amplitudes with permutation operators in front. Thus the two first terms on the right hand side of Eq. (3.18) define $\tilde{V}$ and all the rest $\tilde{R}$. In a standard manner one can sum up $\tilde{V}$ to infinite order leading to

$$\tilde{\psi} = \tilde{\psi}^0 + \tilde{G}_0 \tilde{t} \tilde{R} \quad (3.20)$$

where $\tilde{t}$ obeys the matrix Lippmann-Schwinger equation

$$\tilde{t} = \tilde{V} + \tilde{V} \tilde{G}_0 \tilde{t} \quad (3.21)$$

If one regards a scattering process initiated by some particle hitting a two-baryon bound state then the driving term $\tilde{\psi}^0$ contains a nonzero entry in the corresponding row. For a bound system $\tilde{\psi}^0$ is absent.

As a first example let us regard the strangeness -1 system consisting of the N , Λ and Σ

particles. Let us group the channels in the following manner: ΛNN , ΣNN , $N\Lambda N$, $N\Sigma N$.

Then the matrix $\tilde{V}$ has the form

$$\tilde{V} = \begin{pmatrix} V_{NN} & 0 & 0 & 0 \\ 0 & V_{NN} & 0 & 0 \\ 0 & 0 & V_{\Lambda N, \Lambda N}^{tot} & V_{\Lambda N, \Sigma N}^{tot} \\ 0 & 0 & V_{\Sigma N, \Lambda N}^{tot} & V_{\Sigma N, \Sigma N}^{tot} \end{pmatrix} \quad (3.22)$$

Consequently the Lippman-Schwinger equation (3.21) will decay into two uncoupled equations ($Y = \Lambda$ and Σ)

$$t_{NN}^Y = V_{NN} + N_{NN} G_0^{YNN} t_{NN}^Y \quad (3.23)$$

and one set of two coupled equations

$$\tilde{t} = \begin{pmatrix} V_{\Lambda N, \Lambda N}^{tot} & V_{\Lambda N, \Sigma N}^{tot} \\ V_{\Sigma N, \Lambda N}^{tot} & V_{\Sigma N, \Sigma N}^{tot} \end{pmatrix} + \begin{pmatrix} V_{\Lambda N, \Lambda N}^{tot} & V_{\Lambda N, \Sigma N}^{tot} \\ V_{\Sigma N, \Lambda N}^{tot} & V_{\Sigma N, \Sigma N}^{tot} \end{pmatrix} \begin{pmatrix} G_0^{N\Lambda N} & 0 \\ 0 & G_0^{N\Sigma N} \end{pmatrix} \tilde{t} \quad (3.24)$$

We thus end up with the following four coupled Faddeev equations

$$\psi_{\Lambda NN} = \psi_{\Lambda d}^0 + G_0^{\Lambda NN} t_{NN}^{\Lambda} (1 - P_{23}) P_{13} P_{23} \psi_{\Lambda NN} \quad (3.25)$$

$$\psi_{\Sigma NN} = \psi_{\Sigma d}^0 + G_0^{\Sigma NN} t_{NN}^{\Sigma} (1 - P_{23}) P_{13} P_{23} \psi_{\Sigma NN} \quad (3.26)$$

$$\begin{pmatrix} \psi_{N\Lambda N} \\ \psi_{N\Sigma N} \end{pmatrix} = \begin{pmatrix} G_0^{N\Lambda N} & 0 \\ 0 & G_0^{N\Sigma N} \end{pmatrix} \tilde{t} \begin{pmatrix} -P_{13} \psi_{N\Lambda N} + P_{12} P_{23} \psi_{\Lambda NN} \\ -P_{13} \psi_{N\Sigma N} + P_{12} P_{23} \psi_{\Sigma NN} \end{pmatrix} \quad (3.27)$$

The driving terms in Eqs.(3.25) and (3.26) occur if we regard Λd or Σd scattering. Of course only the one or the other will be nonzero.

It is an easy exercise using appropriate permutations to verify that the homogeneous set corresponding to Eqs. (3.25)-(3.27) is identical to the set solved numerically in [4]. The total hypertriton state Ψ is then given by (3.3). This can be projected into the various particle species.

The case of strangeness $S = -2$ is more complex. We use the following arbitrary but fixed grouping of particle states: ΞNN , $N\Xi N$, $N\Lambda\Sigma$, $N\Lambda\Lambda$, $N\Sigma\Sigma$, $\Lambda N\Lambda$, $\Lambda N\Sigma$, $\Sigma N\Lambda$ and $\Sigma N\Sigma$. Note the remaining six groupings can be related to the nine given ones with the help of Eq. (3.11). Then we can read off from Eq.(3.18) a 9×9 potential matrix. For the sake of an easier notation we drop the Σ particle and find

$$\tilde{V} = \begin{pmatrix} V_{NN} & 0 & 0 & 0 \\ 0 & V_{\Xi N, \Xi N} & V_{\Xi N, \Lambda\Lambda} & 0 \\ 0 & V_{\Lambda\Lambda, \Xi N}^{tot} & V_{\Lambda\Lambda} & 0 \\ 0 & 0 & 0 & V_{N\Lambda, N\Lambda}^{tot} \end{pmatrix} \quad (3.28)$$

Again the matrix Lippman-Schwinger equation for $\tilde{t}$ decays in a corresponding manner and we find the following four coupled Faddeev equations.

$$\psi_{\Xi NN} = \phi_{\Xi d} + G_0^{\Xi NN} t_{NN} (1 - P_{23}) P_{13} P_{23} \psi_{N\Xi N} \quad (3.29)$$

$$\begin{pmatrix} \psi_{N\Xi N} \\ \psi_{N\Lambda\Lambda} \end{pmatrix} = \begin{pmatrix} G_0^{N\Xi N} & 0 \\ 0 & G_0^{N\Lambda\Lambda} \end{pmatrix} \tilde{t} \begin{pmatrix} P_{12} P_{23} \psi_{\Xi NN} - P_{13} \psi_{N\Xi N} \\ (1 - P_{23}) P_{13} P_{23} \psi_{\Lambda N\Lambda} \end{pmatrix} \quad (3.30)$$

$$\psi_{\Lambda N\Lambda} = G_0^{\Lambda N\Lambda} t_{N\Lambda} (P_{12} P_{23} \psi_{N\Lambda\Lambda} - P_{13} \psi_{\Lambda N\Lambda}) \quad (3.31)$$

These four equations are identical to the ones presented in [1]. In order to demonstrate that identity it is again necessary to apply appropriate permutations. This is left to the reader. For later use it is helpful to write that set (3.29)-(3.31) explicitly into the form (3.20). One

easily finds

$$\begin{pmatrix} \psi_{\Xi NN} \\ \psi_{N\Xi N} \\ \psi_{N\Lambda\Lambda} \\ \psi_{\Lambda N\Lambda} \end{pmatrix} = \begin{pmatrix} \phi_{\Xi d} \\ 0 \\ 0 \\ 0 \end{pmatrix}$$

$$+ \begin{pmatrix} G_0^{\Xi NN} & 0 & 0 & 0 \\ 0 & G_0^{N\Xi N} & 0 & 0 \\ 0 & 0 & G_0^{N\Lambda\Lambda} & 0 \\ 0 & 0 & 0 & G_0^{\Lambda N\Lambda} \end{pmatrix} \begin{pmatrix} t_{NN} & 0 & 0 & 0 \\ 0 & t_{\Xi N, \Xi N} & t_{\Xi N, \Lambda\Lambda} & 0 \\ 0 & t_{\Lambda\Lambda, \Xi N} & t_{\Lambda\Lambda, \Lambda\Lambda} & 0 \\ 0 & 0 & 0 & t_{N\Lambda, N\Lambda} \end{pmatrix} \quad (3.32)$$

$$\begin{pmatrix} 0 & (1 - P_{23})P_{13}P_{23} & 0 & 0 \\ P_{12}P_{23} & -P_{13} & 0 & 0 \\ 0 & 0 & 0 & (1 - P_{23})P_{13}P_{23} \\ 0 & 0 & P_{12}P_{23} & -P_{13} \end{pmatrix} \begin{pmatrix} \psi_{\Xi NN} \\ \psi_{N\Xi N} \\ \psi_{N\Lambda\Lambda} \\ \psi_{\Lambda N\Lambda} \end{pmatrix}$$

This identifies explicitly the term $\tilde{t}\tilde{R}$ as $\tilde{t}\tilde{P}\tilde{\psi}$, where the matrix of permutation operators $\tilde{P}$ can be read off from Eq. (3.32). Then Eq.(3.20) can be expressed as

$$\tilde{\psi} = \tilde{\psi}^0 + \tilde{G}_0 \tilde{t} \tilde{P} \tilde{\psi} \quad (3.33)$$

In case of a bound three-body system the driving term $\phi_{\Xi d}$, which is a product of a momentum eigenstate for the Ξ particle and the deuteron state, is of course absent. It is not yet known for sure whether that interesting system fluctuating between ΞNN , $\Lambda\Lambda N$, $\Lambda\Sigma N$ and $\Sigma\Sigma N$ and which is described by a set of nine coupled Faddeev equations is bound or exists at least as a resonance.

IV. BREAKUP PROCESSES IN ΞD SCATTERING

In [2] it was proposed that the process $\Xi d \rightarrow \Lambda\Lambda N$ might be a source of information for the $\Lambda\Lambda$ scattering length. If the $\Lambda\Lambda$ system has a virtual state close to zero relative energy one has to expect an enhancement in the breakup configuration, when the two Λ 's leave with equal laboratory momenta. In [2] it was found in a numerical study using model forces that a destructive interference between different production amplitudes diminishes that enhancement. The strength of that suppression turned out to be linked to the $\Lambda\Lambda - \Xi N$ coupling and is therefore of great theoretical interest. As already touched upon in the introduction the formalism in [2] is based on the AGS equations for distinguishable particles and antisymmetrisation is attached through the antisymmetrised asymptotic initial and final channel states. Obviously it is desirable to have a straightforward derivation of the appropriate scattering formalism. This is achieved in the form of the coupled set of Faddeev equations (3.29)-(3.31) for the four independent Faddeev amplitudes. (In case of an additional Σ -admixture one has nine independent amplitudes.) They describe the elastic process

$$\Xi + d \rightarrow \Xi + d \tag{4.1}$$

and the various breakup processes

$$\Xi + d \rightarrow \begin{cases} \Xi + N + N \\ \Lambda + \Lambda + N \\ N + \Lambda + \Sigma \\ N + \Sigma + \Sigma \end{cases} \tag{4.2}$$

It is a standard procedure [9–12] to extract the various partial breakup amplitudes from each individual kernel part of the set (3.29)-(3.31) or in general matrix notation from the kernel of (3.20). One regards the asymptotic behavior in configuration space and finds the

partial breakup amplitudes to be

$$\tilde{T} \equiv \tilde{t}\tilde{R} \tag{4.3}$$

Since $\tilde{R} = \tilde{P}\tilde{\psi}$ and

$$\tilde{\psi} = \tilde{\psi}^0 + \tilde{G}_0\tilde{T} \tag{4.4}$$

one arrives at

$$\tilde{T} = \tilde{t}\tilde{P}\tilde{\psi}^0 + \tilde{t}\tilde{P}\tilde{G}_0\tilde{T} \tag{4.5}$$

which is a closed set of equations for all partial breakup amplitudes. Again for the sake of a simple notation we regard the set (3.29)-(3.31) combined in Eq.(3.32) without Σ and easily find that Eq. (4.5) takes the explicit form:

$$\begin{pmatrix} T_{\Xi NN} \\ T_{N\Xi N} \\ T_{N\Lambda\Lambda} \\ T_{\Lambda N\Lambda} \end{pmatrix} = \begin{pmatrix} 0 \\ t_{\Xi N, \Xi N} P_{12} P_{23} \phi_{\Xi d} \\ t_{\Lambda\Lambda, \Xi N} P_{12} P_{23} \phi_{\Xi d} \\ 0 \end{pmatrix}$$

$$+ \begin{pmatrix} 0 & t_{NN}(1 - P_{23})P_{13}P_{23} & 0 & 0 \\ t_{\Xi N, \Xi N} P_{12} P_{23} & -t_{\Xi N, \Xi N} P_{13} & 0 & t_{\Xi N, \Lambda\Lambda}(1 - P_{23})P_{13}P_{23} \\ t_{\Lambda\Lambda, \Xi N} P_{12} P_{23} & -t_{\Lambda\Lambda, \Xi N} P_{13} & 0 & t_{\Lambda\Lambda, \Lambda\Lambda}(1 - P_{23})P_{13}P_{23} \\ 0 & 0 & t_{N\Lambda, N\Lambda} P_{12} P_{23} & -t_{N\Lambda, N\Lambda} P_{13} \end{pmatrix} \quad (4.6)$$

$$\begin{pmatrix} G_0^{\Xi NN} & 0 & 0 & 0 \\ 0 & G_0^{N\Xi N} & 0 & 0 \\ 0 & 0 & G_0^{N\Lambda\Lambda} & 0 \\ 0 & 0 & 0 & G_0^{\Lambda N\Lambda} \end{pmatrix} \begin{pmatrix} T_{\Xi NN} \\ T_{N\Xi N} \\ T_{N\Lambda\Lambda} \\ T_{\Lambda N\Lambda} \end{pmatrix}$$

The full breakup amplitudes result from projecting the full wavefunction given in Eq. (3.3) onto the various breakup configurations. Using (3.15) one finds for instance

$$\langle \Xi NN | \Psi \rangle = \langle \Xi NN | (1 + P) | \psi \rangle = \psi_{\Xi NN} + (1 - P_{23})P_{13}P_{23}\psi_{N\Xi N} \quad (4.7)$$

Consequently the full breakup amplitude into the channel ΞNN is

$$U_{\Xi NN}^0 = T_{\Xi NN} + (1 - P_{23})P_{13}P_{23}T_{N\Xi N} \quad (4.8)$$

Similarly we get for the breakup process into $\Lambda N\Lambda$

$$U_{\Lambda N\Lambda}^0 = (1 - P_{13})T_{\Lambda N\Lambda} + P_{12}P_{23}T_{N\Lambda\Lambda} \quad (4.9)$$

Note both amplitudes have the appropriate antisymmetry for NN and $\Lambda\Lambda$, respectively

as guaranteed by the properties $P_{23}T_{\Xi NN} = -T_{\Xi NN}$ and $P_{23}T_{N\Lambda\Lambda} = -T_{N\Lambda\Lambda}$. All further projections lead to no new breakup amplitudes. If we would have projected for instance in case that in addition the Σ is included for the $\Lambda\Sigma N$ breakup the resulting amplitude $\langle \Lambda\Sigma N | \Psi \rangle$ is connected to $\langle N\Lambda\Sigma | \Psi \rangle$ by $P_{23}P_{13} \langle \Lambda\Sigma N | \Psi \rangle = \langle N\Lambda\Sigma | \Psi \rangle$ which tells that only the particles have been numbered differently.

Since after all those YN and YY forces are weaker than the NN force it is not unreasonable to look into a multiple scattering series expansion of the set (4.5). It might even converge if in the future it will turn out that there is no strangeness -2 three-baryon bound state nor even a low lying three-body resonance. The multiple scattering series results by iterating the set(4.6). In lowest and next to lowest order in the t -operators one gets

$$T_{N\Xi N} = -t_{\Xi N, \Xi N}(23)P_{12}\phi_{\Xi d} + t_{\Xi N, \Xi N}(23)P_{13}G_0^{N\Xi N}t_{\Xi N, \Xi N}(23)P_{12}\phi_{\Xi d} \quad (4.10)$$

$$T_{N\Lambda\Lambda} = -t_{\Lambda\Lambda, \Xi N}(23)P_{12}\phi_{\Xi d} + t_{\Lambda\Lambda, \Xi N}(23)P_{13}G_0^{N\Xi N}t_{\Xi N, \Xi N}(23)P_{12}\phi_{\Xi d} \quad (4.11)$$

$$T_{\Lambda N\Lambda} = -t_{N\Lambda, N\Lambda}(23)P_{12}P_{23}G_0^{N\Lambda\Lambda}t_{\Lambda\Lambda, \Xi N}(23)P_{12}\phi_{\Xi d} \quad (4.12)$$

$$T_{\Xi NN} = t_{NN}(23)(1 - P_{23})P_{12}G_0^{N\Xi N}t_{\Xi N, \Xi N}(23)P_{12}\phi_{\Xi d} \quad (4.13)$$

Using Eqs. (4.8) and (4.9) we arrive at the two breakup amplitudes to second order in the t -matrices:

$$\begin{aligned} U_{\Xi NN}^0 &= t_{\Xi N, \Xi N}(12)\phi_{\Xi d} + t_{\Xi N, \Xi N}(13)\phi_{\Xi d} \\ &+ t_{\Xi N, \Xi N}(12)G_0^{\Xi NN}t_{\Xi N, \Xi N}(13)\phi_{\Xi d} + t_{\Xi N, \Xi N}(13)G_0^{\Xi NN}t_{\Xi N, \Xi N}(12)\phi_{\Xi d} \\ &+ t_{NN}(23)G_0^{\Xi NN}t_{\Xi N, \Xi N}(13)\phi_{\Xi d} + t_{NN}(23)G_0^{\Xi NN}t_{\Xi N, \Xi N}(12)\phi_{\Xi d} \end{aligned} \quad (4.14)$$

$$\begin{aligned}
U_{\Lambda N \Lambda}^0 &= t_{\Lambda \Lambda, \Xi N}(13) \phi_{\Xi d} \\
&+ t_{N \Lambda, N \Lambda}(23) G_0^{\Lambda N \Lambda} t_{\Lambda \Lambda, \Xi N}(13) \phi_{\Xi d} + t_{N \Lambda, N \Lambda}(21) G_0^{\Lambda N \Lambda} t_{\Lambda \Lambda, \Xi N}(13) \phi_{\Xi d} \\
&+ t_{\Lambda \Lambda, \Xi N}(13) G_0^{\Xi N N} t_{\Xi N, \Xi N}(12) \phi_{\Xi d}
\end{aligned} \tag{4.15}$$

In arriving at Eq. (4.15) we used

$$P_{23} t_{\Lambda \Lambda, \Xi N}(23) = -t_{\Lambda \Lambda, \Xi N}(23) \tag{4.16}$$

which is direct consequence of the definition (2.16) of $V_{\alpha\beta, \alpha'\beta'}^{tot}$ and the corresponding Lippmann-Schwinger equation (see Appendix). The antisymmetry of $U_{\Xi N N}^0$ in the two nucleons, $P_{23} U_{\Xi N N}^0 = -U_{\Xi N N}^0$, can easily be seen to be guaranteed by the antisymmetry of the deuteron state. In case of $U_{\Lambda N \Lambda}^0$ the antisymmetry $P_{13} U_{\Lambda N \Lambda}^0 = -U_{\Lambda N \Lambda}^0$ is due to the property (4.16). We exhibit the content of Eqs. (4.14) and (4.15) graphically in Figs. 1 and 2. With the techniques available nowadays (see for instance [11]) it is feasible to evaluate these processes reliably using modern forces.

V. ELASTIC ΞD SCATTERING-GENERALIZED AGS EQUATIONS

The two sets of four coupled Faddeev equations (3.25-3.27) and (3.29)-(3.31) or more generally (3.33) also define the asymptotic behavior in bound state channels. As known up to now there is only one baryon-baryon bound state, the deuteron. Using the well known relation

$$(E - H_0)^{-1} t_{NN} = (E - H_0 - V_{NN})^{-1} V_{NN} \tag{5.1}$$

one can extract easily the asymptotic behavior in configuration space in the deuteron channel [12]. For instance the elastic amplitude in Ξd scattering extracted from Eq. (3.29) is

$$U_{\Xi d} = \langle \phi'_{\Xi d} | V_{NN}(23)(1 - P_{23})P_{13}P_{23} | \psi_{N\Xi N} \rangle \quad (5.2)$$

where the prime on the channel state $\phi'_{\Xi d}$ indicates that the momentum of the outgoing Ξ -particle is independent of the momentum of the incoming Ξ -particle. Inserting the right hand side of the Faddeev equation from Eq. (3.30) for $\psi_{N\Xi N}$ and using the definition of $T_{N\Xi N}$ as expressed in Eq. (4.4) one finds

$$U_{\Xi d} = \langle \phi'_{\Xi d} | V_{NN}(23)(1 - P_{23})P_{13}P_{23}G_0^{N\Xi N}T_{N\Xi N} \rangle \quad (5.3)$$

Thus the elastic amplitude can be determined by quadrature from the partial breakup amplitude $T_{N\Xi N}$.

Though for practical calculations working with general forces the Faddeev equation for the T 's are most appropriate [11] it is interesting from a formal point of view to derive Faddeev equations for transition operators describing elastic and rearrangement scattering. Using forces of finite rank they are also a good starting point.

In the case of the outgoing Ξd channel we extracted a state

$$U'_{\Xi d \Xi d} \equiv V_{NN}(23)(1 - P_{23})P_{13}P_{23}\psi_{N\Xi N} \quad (5.4)$$

which under action from the left by $\langle \phi'_{\Xi d} |$ gives the elastic transition amplitude (see Eq. (5.2)). In the elastic amplitude on the energy shell we apparently can replace V_{NN} by $(G_0^{\Xi NN})^{-1} = (E - H_0^{\Xi NN})$. Let us therefore define

$$U_{\Xi d \Xi d} \equiv (G_0^{\Xi NN})^{-1}(1 - P_{23})P_{13}P_{23}\psi_{N\Xi N} \quad (5.5)$$

Analogously we define

$$\begin{pmatrix} U_{N\Xi N, \Xi d} \\ U_{N\Lambda\Lambda, \Xi d} \end{pmatrix} \equiv \begin{pmatrix} (G_0^{N\Xi N})^{-1} & 0 \\ 0 & (G_0^{N\Lambda\Lambda})^{-1} \end{pmatrix} \begin{pmatrix} P_{12}P_{23}\psi_{\Xi NN} - P_{13}\psi_{N\Xi N} \\ (1 - P_{23})P_{13}P_{23}\psi_{\Lambda N\Lambda} \end{pmatrix} \quad (5.6)$$

$$U_{\Lambda N \Lambda, \Xi d} \equiv (G_0^{\Lambda N \Lambda})^{-1} (P_{12} P_{23} \psi_{N \Lambda \Lambda} - P_{13} \psi_{\Lambda N \Lambda}) \quad (5.7)$$

These definitions together with the coupled set (3.29)-(3.31) will lead to a closed set of equations for the four quantities U . It is advisable to write the four defining equations (5.5)-(5.7) into a matrix form

$$\begin{aligned} \tilde{U} \equiv \begin{pmatrix} U_{\Xi d, \Xi d} \\ U_{N \Xi N, \Xi d} \\ U_{N \Lambda \Lambda, \Xi d} \\ U_{\Lambda N \Lambda, \Xi d} \end{pmatrix} &= \begin{pmatrix} (G_0^{\Xi N N})^{-1} & 0 & 0 & 0 \\ 0 & (G_0^{N \Xi N})^{-1} & 0 & 0 \\ 0 & 0 & (G_0^{N \Lambda \Lambda})^{-1} & 0 \\ 0 & 0 & 0 & (G_0^{\Lambda N \Lambda})^{-1} \end{pmatrix} \\ &\begin{pmatrix} 0 & (1 - P_{23}) P_{13} P_{23} & 0 & 0 \\ P_{12} P_{23} & -P_{13} & 0 & 0 \\ 0 & 0 & 0 & (1 - P_{23}) P_{13} P_{23} \\ 0 & 0 & P_{12} P_{23} & -P_{13} \end{pmatrix} \begin{pmatrix} \psi_{\Xi N N} \\ \psi_{N \Xi N} \\ \psi_{N \Lambda \Lambda} \\ \psi_{\Lambda N \Lambda} \end{pmatrix} \quad (5.8) \\ &\equiv \tilde{G}_0^{-1} \tilde{P} \tilde{\psi} \end{aligned}$$

We recover the matrix $\tilde{P}$ of permutations already occurring in Eq. (3.32) and in the coupled set (3.33).

Generally spoken that matrix Faddeev equation (3.33) is a special case of Eq. (3.20) where

$$\tilde{R} \equiv \tilde{P} \tilde{\psi} \quad (5.9)$$

contains now a general matrix $\tilde{P}$ of permutations (see below). Now we can proceed and insert $\tilde{\psi}$ from (3.20) together with (5.9) back into (5.8) and find

$$\tilde{U} = \tilde{G}_0^{-1} \tilde{P} (\tilde{\psi}^0 + \tilde{G}_0 \tilde{t} \tilde{P} \tilde{\psi}) = \tilde{G}_0^{-1} \tilde{P} \tilde{\psi}^0 + \tilde{G}_0^{-1} \tilde{P} \tilde{G}_0 \tilde{t} \tilde{G}_0 \tilde{U} \quad (5.10)$$

It remains to demonstrate that

$$\tilde{G}_0^{-1}\tilde{P} = \tilde{P}\tilde{G}_0^{-1} \quad (5.11)$$

and thus

$$\tilde{G}_0^{-1}\tilde{P}\tilde{G}_0 = \tilde{P} \quad (5.12)$$

Written out explicitly we have

$$\tilde{P}\tilde{G}_0^{-1} = \begin{pmatrix} 0 & (1 - P_{23})P_{13}P_{23}(E - H_0^{N\Xi N}) & 0 & 0 \\ P_{12}P_{23}(E - H_0^{\Xi NN}) & -P_{13}(E - H_0^{N\Xi N}) & 0 & 0 \\ 0 & 0 & 0 & (1 - P_{23})P_{13}P_{23}(E - H_0^{\Lambda N\Lambda}) \\ 0 & 0 & P_{12}P_{23}(E - H_0^{N\Lambda\Lambda}) & -P_{13}(E - H_0^{\Lambda N\Lambda}) \end{pmatrix} \quad (5.13)$$

Let us regard two examples

$$\begin{aligned} & (1 - P_{23})P_{13}P_{23}H_0^{N\Xi N}(123) \\ &= (1 - P_{23})H_0^{N\Xi N}(312)P_{13}P_{23} \\ &= (1 - P_{23})H_0^{\Xi NN}(123)P_{13}P_{23} \\ &= H_0^{\Xi NN}(123)(1 - P_{23})P_{13}P_{23} \end{aligned} \quad (5.14)$$

$$P_{12}P_{23}H_0^{\Xi NN}(123) = H_0^{\Xi NN}(231)P_{12}P_{23} = H_0^{N\Xi N}(123)P_{12}P_{23} \quad (5.15)$$

Corresponding relations are true in all cases and we find (5.12) to be valid. Thus we end up with the coupled matrix Faddeev equation

$$\tilde{U} = \tilde{P}\tilde{G}_0^{-1}\tilde{\psi}^0 + \tilde{P}\tilde{t}\tilde{G}_0\tilde{U} \quad (5.16)$$

For the case of the coupled $\Xi NN - \Lambda\Lambda N$ system they read in explicit form

$$U_{\Xi d \Xi d} = (1 - P_{23})P_{13}P_{23}(t_{\Xi N, \Xi N}G_0^{N\Xi N}U_{N\Xi N, \Xi d} + t_{\Xi N, \Lambda\Lambda}G_0^{N\Lambda\Lambda}U_{N\Lambda\Lambda, \Xi d}) \quad (5.17)$$

$$\begin{aligned} U_{N\Xi N, \Xi d} &= P_{12}P_{23}(G_0^{\Xi NN})^{-1}\phi_{\Xi d} \\ &+ P_{12}P_{23}t_{NN}G_0^{\Xi NN}U_{\Xi d \Xi d} - P_{13}(t_{\Xi N, \Xi N}G_0^{N\Xi N}U_{N\Xi N, \Xi d} + t_{\Xi N, \Lambda\Lambda}G_0^{N\Lambda\Lambda}U_{N\Lambda\Lambda, \Xi d}) \end{aligned} \quad (5.18)$$

$$U_{N\Lambda\Lambda, \Xi d} = (1 - P_{23})P_{13}P_{23}t_{N\Lambda, N\Lambda}G_0^{\Lambda N\Lambda}U_{\Lambda N\Lambda, \Xi d} \quad (5.19)$$

$$U_{\Lambda N\Lambda, \Xi d} = P_{12}P_{23}(t_{\Lambda\Lambda, \Xi N}G_0^{N\Xi N}U_{N\Xi N, \Xi d} + t_{\Lambda\Lambda, \Lambda\Lambda}G_0^{N\Lambda\Lambda}U_{N\Lambda\Lambda, \Xi d}) - P_{13}t_{N\Lambda, N\Lambda}G_0^{\Lambda N\Lambda}U_{\Lambda N\Lambda, \Xi d} \quad (5.20)$$

This is a generalized set of Alt-Grassberger-Sandhas (AGS) equations for that specific system. Obviously a corresponding set can be written down for the coupled $\Lambda NN - \Sigma NN$ system based on the four Faddeev equations (3.25-3.27).

The matrix equation (5.16) (generalized AGS equations) is generally valid. The proof is quite simple. In the set (3.19) the part including permutations is by definition

$$\langle \alpha\beta\gamma | G_0 V P | \psi \rangle = \sum_{\alpha'\beta'\gamma'} \sum_{\alpha''\beta''\gamma''} G_0^{\alpha\beta\gamma} \langle \alpha\beta\gamma | V | \alpha'\beta'\gamma' \rangle \langle \alpha'\beta'\gamma' | P | \alpha''\beta''\gamma'' \rangle \psi_{\alpha''\beta''\gamma''} \quad (5.21)$$

This defines the general matrix for the permutation operators. Now $\tilde{G}_0\tilde{P}$ contains the matrix element

$$G_0^{\alpha\beta\gamma} \langle \alpha\beta\gamma | P | \alpha'\beta'\gamma' \rangle = \langle \alpha\beta\gamma | G_0 P | \alpha'\beta'\gamma' \rangle \quad (5.22)$$

The operator $G_0 \equiv (E - H_0)^{-1}$ occurring in Eqs. (3.4-3.5) is however fully symmetrical.

Thus $G_0P = PG_0$ and the relation (5.12) follows.

Finally we note that due to (5.8) and (4.4) the matrix $\tilde{U}$ of elastic and rearrangement quantities is related to the matrix $\tilde{T}$ composed of partial breakup amplitudes via

$$\tilde{U} = \tilde{P}G_0^{-1}\tilde{\psi}^0 + \tilde{P}\tilde{T} \quad (5.23)$$

and can therefore be determined by quadrature once $\tilde{T}$ is evaluated from Eq. (4.5). We remark that in the 3N system (three identical particles) all those matrix relations reduce to the well known single equations [11]:

$$U = PG_0^{-1}\psi^0 + PT \quad (5.24)$$

$$T = tP\psi^0 + tPG_0T \quad (5.25)$$

$$U = PG_0^{-1}\psi^0 + PtG_0U \quad (5.26)$$

In that form they have been solved rigorously (in a numerical sense) for modern nuclear forces.

VI. SUMMARY

A symmetric Hamiltonian for distinguishable particles with different masses and non-symmetric interactions has been constructed using the generalized Pauli principle based on particle labels added to the usual ones for space (momentum) coordinates and spin. This has been exemplified in two-body systems and then applied to three-baryon systems. Specifically we had in mind the strangeness $S = -1$ and -2 systems. We derived minimal sets of coupled Faddeev equations for bound states, for breakup operators and rearrangement

operators (generalized AGS equations). In the case of the coupled $\Lambda NN - \Sigma NN$ system we arrive at four Faddeev equations. Dropping a possible Σ -admixture the system $\Xi NN - \Lambda\Lambda N$ is also described by four Faddeev equations. Allowing for a Σ -admixture one would have nine equations. An application of such a set of equations already appeared in case of ${}^3_{\Lambda}\text{H}$ in [4]. An investigation of the $S = -2$ system is planned.

ACKNOWLEDGMENTS

We thank M.Oka for a stimulating discussion. This work was performed during visits of W.G. at the Okayama University of Science and of K.M. at the Ruhr-University Bochum. Both authors thank for the hospitality extended to them at the two institutions.

REFERENCES

- [1] K. Miyagawa, H. Kamada, and W. Glöckle, Nucl. Phys. A614, 535 (1997).
- [2] S.B.Carr, I.R.Afnan, B.F.Gibson, Phys.Rev. C57(1998) 2858.
- [3] E.O.Alt, P.Grassberger, W.Sandhas, Nucl. Phys. B2 (1967) 167
- [4] K.Miyagawa, H.Kamada, W.Glöckle, V.Stoks, Phys Rev. C51, 2905(1995).
- [5] E.Hiyama, in Proceedings of the XVII RCNP international symposium on innovative computational method in nuclear many-body problems, Osaka, November 1997, eds. H.Horiuchi, M.Kamimura, H.Toki, Y.Fujiwara, M.Matsuo, Y.Sakuragi, World Scientific 1998, page 128.
- [6] H.Yamamura, K.Miyagawa, T.Mart, C.Bennhold, H.Haberzettl, W.Glöckle, Phys. Rev. C61, 014001-1(1999).
- [7] D.M.Koltenuk, PhD thesis, University of Pennsylvania, 1999, unpublished.
- [8] T.-S.H. Lee, V.Stoks, B.Saghai, C.Fayard, Nucl. Phys. A639, 247c (1998).
- [9] W.Glöckle, The Quantum Mechanical Few-Body Problem, Springer Verlag, 1983.
- [10] D.Hüber, H.Kamada, H.Witala, W.Glöckle, Acta Phys. Polonica B21, 1677(1997).
- [11] W.Glöckle, H.Witala, D.Hüber, H.Kamada, J.Golak, Phys. Rep. 274, 107(1996).
- [12] W.Glöckle, Zeitschrift f.Phys. 271, 31(1974).

Appendix: The property (4.16)

Starting from the Lippman-Schwinger equation

$$\begin{pmatrix} t_{\Xi N, \Xi N} & t_{\Xi N, \Lambda\Lambda} \\ t_{\Lambda\Lambda, \Xi N} & t_{\Lambda\Lambda, \Lambda\Lambda} \end{pmatrix} = \begin{pmatrix} V_{\Xi N, \Xi N} & V_{\Xi N, \Lambda\Lambda} \\ V_{\Lambda\Lambda, \Xi N}^{tot} & V_{\Lambda\Lambda, \Lambda\Lambda} \end{pmatrix} + \begin{pmatrix} V_{\Xi N, \Xi N} & V_{\Xi N, \Lambda\Lambda} \\ V_{\Lambda\Lambda, \Xi N}^{tot} & V_{\Lambda\Lambda, \Lambda\Lambda} \end{pmatrix} \begin{pmatrix} G_0^{\Xi NN} & 0 \\ 0 & G_0^{\Lambda\Lambda N} \end{pmatrix} \begin{pmatrix} t_{\Xi N, \Xi N} & t_{\Xi N, \Lambda\Lambda} \\ t_{\Lambda\Lambda, \Xi N} & t_{\Lambda\Lambda, \Lambda\Lambda} \end{pmatrix} \quad (1)$$

one finds

$$t_{\Lambda\Lambda, \Xi N} = V_{\Lambda\Lambda, \Xi N}^{tot} + V_{\Lambda\Lambda, \Xi N}^{tot} G_0^{\Xi NN} t_{\Xi N, \Xi N} + V_{\Lambda\Lambda, \Lambda\Lambda} G_0^{\Lambda\Lambda N} t_{\Lambda\Lambda, \Xi N} \quad (2)$$

Here we number the interacting particles as 1 and 2. Applying P_{12} yields

$$\begin{aligned} P_{12} t_{\Lambda\Lambda, \Xi N} &= P_{12} V_{\Lambda\Lambda, \Xi N}^{tot} + P_{12} V_{\Lambda\Lambda, \Xi N}^{tot} G_0^{\Xi NN} t_{\Xi N, \Xi N} \\ &\quad + P_{12} V_{\Lambda\Lambda, \Lambda\Lambda} G_0^{\Lambda\Lambda N} t_{\Lambda\Lambda, \Xi N} \\ &= -V_{\Lambda\Lambda, \Xi N}^{tot} - V_{\Lambda\Lambda, \Xi N}^{tot} G_0^{\Xi NN} t_{\Xi N, \Xi N} + V_{\Lambda\Lambda, \Lambda\Lambda} G_0^{\Lambda\Lambda N} P_{12} t_{\Lambda\Lambda, \Xi N} \end{aligned} \quad (3)$$

We used the definition (2.16) and the obvious symmetry of $V_{\Lambda\Lambda, \Lambda\Lambda} G_0^{\Lambda\Lambda N}$ under exchange of particles 1 and 2. It follows

$$\begin{aligned} P_{12} t_{\Lambda\Lambda, \Xi N} &= -(1 - V_{\Lambda\Lambda, \Lambda\Lambda} G_0^{\Lambda\Lambda N})^{-1} (V_{\Lambda\Lambda, \Xi N}^{tot} + V_{\Lambda\Lambda, \Xi N}^{tot} G_0^{\Xi NN} t_{\Xi N, \Xi N}) \\ &= -t_{\Lambda\Lambda, \Xi N} \end{aligned} \quad (4)$$

which is claimed in (4.16).

FIGURES




FIG. 1. The ΞNN breakup process of Eq. (4.14) up to second order in the t -matrices. The hatched half circle denotes the incoming deuteron.




FIG. 2. The $\Lambda N \Lambda$ breakup process of Eq. (4.15) up to second order in the t -matrices.